

Block-Spiral Magnetism: A New Type of Frustrated Order

J. Herbrych^{1,2,3}, J. Heverhagen^{4,5}, G. Alvarez⁶, M. Daghofer^{4,5}, A. Moreo^{1,2}, and E. Dagotto^{1,2}

¹ Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA

² Materials Science and Technology Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

³ Department of Theoretical Physics, Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology, 50-370 Wrocław, Poland

⁴ Institute for Functional Matter and Quantum Technologies, University of Stuttgart, Pfaffenwaldring 57, D-70550 Stuttgart, Germany

⁵ Center for Integrated Quantum Science and Technology, University of Stuttgart, Pfaffenwaldring 57, D-70550 Stuttgart, Germany and

⁶ Computational Sciences and Engineering Division and Center for Nanophase Materials Sciences, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

(Dated: November 28, 2019)

Competing interactions in Quantum Materials induce novel states of matter such as frustrated magnets, an extensive field of research both from the theoretical and experimental perspectives. Here, we show that competing energy scales present in the low-dimensional orbital-selective Mott phase (OSMP) induce an exotic magnetic order, never reported before. Earlier neutron scattering experiments on iron-based 123 ladder materials, where OSMP is relevant, already confirmed our previous theoretical prediction of block-magnetism (magnetic order of the form $\uparrow\uparrow\downarrow\downarrow$). Now we argue that another novel phase can be stabilized in multiorbital Hubbard models, the “block-spiral state”. In this state, the magnetic islands form a spiral propagating through the chain but with the blocks maintaining their identity, namely rigidly rotating. This new spiral state is stabilized without any apparent frustration, the common avenue to generate spiral arrangements in multiferroics. By examining the behaviour of the electronic degrees of freedom, parity breaking quasiparticles are revealed. Finally, a simple phenomenological model that accurately captures the macroscopic spin spiral arrangement is also introduced.

Frustrated magnetism is one of the main areas of research in contemporary Condensed Matter Physics. In the generic scenario, magnetic frustration emerges from the failure of the system to fulfill simultaneously conflicting local requirements. As a consequence, complex spin patterns can develop from geometrical frustration (as in triangular, Kagome, or pyrochlore lattices) or from special spin-spin interactions (long-range exchange, Dzyaloshinskii-Moriya coupling, Kitaev model, spin-orbit effects, and others). In real materials both scenarios often coexist. Competing mechanisms can lead to interesting phenomena, such as spiral order [1–3], spin ice [4], skyrmions [5], spin liquids [6, 7], and resonating valence bond states [8]. Also, the electronic properties of such systems are of much interest: it was shown that the interplay of a spiral state on a metallic host can support Majorana fermions [9–14] and can also induce multiferoicity [15–21].

Another example of competing interactions is the orbital-selective Mott phase (OSMP) [22, 23]. In multi-orbital systems, this effect can lead to the selective localization of electrons on some orbitals. The latter coexist with itinerant bands of mobile electrons. This unique mixture of localized and itinerant components in multi-orbital systems could be responsible for the (bad) metallic behaviour of the parent compounds of iron-based superconductors. This is in strike contrast to cuprates, usually described by the single-band Hubbard model, where parent materials are insulators. Furthermore, the OSMP can also host exotic magnetic phases. It was shown [24]

that the competition between Hund and Hubbard interaction can stabilize unexpected block-magnetism [25–28], namely antiferromagnetically (AFM) coupled ferromagnetic (FM) islands. The size and shape of such islands depends on the electronic filling of the itinerant orbitals. This work focuses on two representative cases: (i) the $\pi/2$ -block spin-pattern $\uparrow\uparrow\downarrow\downarrow$ and (ii) the $\pi/3$ -block spin-pattern $\uparrow\uparrow\downarrow\downarrow\downarrow$. Note that these block-patterns are deduced from the spin-spin correlations vs distance, and not from the local expectation value of the spin operator; these are not spin-density waves.

The OSMP was shown [22] to be relevant for the low-dimensional family of iron-based ladders, the so-called 123 family AFe_2X_3 , where $A=Ba, K, Rb$ is an alkaline earth metal and $X=S, Se$ is a chalcogen. From the magnetism perspective, two phases were experimentally reported: (i) for $BaFe_2Se_3$ inelastic neutron scattering (INS) identified [29, 30] a 2×2 block-magnetic phase in a $\uparrow\uparrow\downarrow\downarrow$ -pattern along the legs. (ii) On the other hand, for $BaFe_2S_3$ and $RbFe_2Se_3$ INS reported [31–33] 2×1 blocks, FM along rungs and AFM along legs. Both of these phases are captured by the multi-orbital Hubbard Hamiltonian [28] and also by its low-energy effective description [24], the generalized Kondo-Heisenberg model.

In this work, we will show that another novel magnetic phase can be stabilized, namely we will report an exotic *block-spiral* state. Such a state arises as a consequence of simultaneous tendencies in the system to form magnetic blocks and to develop non-collinear order. Different from “standard” spirals where one can observe spin-to-

spin rotation, in the block-spiral state the blocks rigidly rotate, as illustrated in the three panels of Fig. 1(a)]. Furthermore, such a magnetic pattern appears without any apparent frustration in the model and is a consequence of subtly competing energy scales present in the OSMP. This novel spin order originates in ferromagnetic tendencies, induced by Hund physics in multi-orbital systems, and opposite antiferromagnetic superexchange tendencies, caused by Hubbard interactions. We propose two simpler phenomenological models that capture the essence of our findings. Moreover, the electronic properties can be accurately described by quasiparticles that break parity symmetry, as expected within a spiral state. Also, we will show that the behaviour of spins can be modeled via a frustrated long-range spin-only Heisenberg model. Such a spin model can serve as a starting point for the spin-wave theory calculations often used to compare theory vs INS spectra. Our findings are robust against system parameter changes and should characterize generic multi-orbital systems in the OSMP regime, close to ferromagnetism.

Results

System Hamiltonians. We will discuss the properties of a multi-orbital Hubbard model on a one-dimensional (1D) lattice. In the generic $SU(2)$ -symmetric form, the two-orbital Hubbard model can be written as

$$\begin{aligned}
 H_H = & - \sum_{\gamma, \gamma', \ell, \sigma} t_{\gamma\gamma'} \left(c_{\gamma, \ell, \sigma}^\dagger c_{\gamma', \ell+1, \sigma} + \text{H.c.} \right) + \Delta \sum_{\ell} n_{1, \ell} \\
 & + U \sum_{\gamma, \ell} n_{\gamma, \ell, \uparrow} n_{\gamma, \ell, \downarrow} + (U - 5J_H/2) \sum_{\ell} n_{0, \ell} n_{1, \ell} \\
 & - 2J_H \sum_{\ell} \mathbf{S}_{0, \ell} \cdot \mathbf{S}_{1, \ell} + J_H \sum_{\ell} \left(P_{0, \ell}^\dagger P_{1, \ell} + \text{H.c.} \right). \quad (1)
 \end{aligned}$$

Here, $c_{\gamma, \ell, \sigma}^\dagger$ ($c_{\gamma, \ell, \sigma}$) creates (destroys) an electron with spin projection $\sigma = \{\uparrow, \downarrow\}$ at orbital $\gamma = \{0, 1\}$ of site $\ell = \{1, \dots, L\}$. Δ stands for crystal-field splitting, while $n_{\gamma, \ell} = \sum_{\sigma} n_{\gamma, \ell, \sigma}$ represents the total density of electrons at (γ, ℓ) , with $n_{\gamma, \ell, \sigma}$ the density of electrons with σ -spin projection. U is the standard, same-orbital repulsive Hubbard interaction, and J_H is the Hund exchange between spins $\mathbf{S}_{\gamma, \ell}$ at different orbitals γ (fixed here to $J_H = U/4$). Finally, the last term $P_{0, \ell}^\dagger P_{1, \ell}$ stands for pair-hopping between orbitals, where $P_{\gamma, \ell} = c_{\gamma, \ell, \uparrow} c_{\gamma, \ell, \downarrow}$.

The choice of hopping matrix elements $t_{\gamma\gamma'}$ – specifically $t_{00} = 0.5$ [eV], $t_{11} = 0.15$ [eV], and $t_{01} = t_{10} = 0$ – and crystal-field splitting $\Delta = 0.8$ [eV] is motivated by our previous studies [24, 28, 34] on the magnetic properties of the OSMP. They represent a generic case of coexisting wide and narrow electronic bands, as often found in iron-based materials from the 123 family [23, 28, 35, 36]. The above values yield a kinetic energy bandwidth $W = 2.1$ eV which is used as an energy unit throughout the paper.




Figure 1. Schematic representation of spirals within the orbital-selective Mott phase. (a) Top to bottom: “standard” spiral spin structure with site-to-site spins rotation, novel block-spirals of two and three sites, respectively. (b) Generalized Kondo-Heisenberg model on a one-dimensional lattice. (c) Interaction U –filling n_H/n_K phase diagram. Solid (dashed) coloring represents the OSMP (paramagnetic) region.

Furthermore, it was previously shown [24] that the magnetic properties of the OSMP can be qualitatively, and even quantitatively, described by the effective Hamiltonian obtained by Schrieffer-Wolff transforming the subspace with strictly one electron per site at the localized orbital $\gamma = 1$, leading to the generalized Kondo-Heisenberg (gKH) model [see also Fig. 1(b)]

$$\begin{aligned}
 H_K = & t_{00} \sum_{\ell, \sigma} \left(c_{0, \ell, \sigma}^\dagger c_{0, \ell+1, \sigma} + \text{H.c.} \right) + U \sum_{\ell} n_{0, \ell, \uparrow} n_{0, \ell, \downarrow} \\
 & + K \sum_{\ell} \mathbf{S}_{1, \ell} \cdot \mathbf{S}_{1, \ell+1} - 2J_H \sum_{\ell} \mathbf{S}_{0, \ell} \cdot \mathbf{S}_{1, \ell}, \quad (2)
 \end{aligned}$$

where $K = 4t_{11}^2/U$. It is worth noting that due to particle-hole symmetry, the electronic filling relation between the two-orbital Hubbard (n_H) and gKH (n_K) mod-




Figure 2. Interaction U/W dependence of correlation functions. (a,d) Static spin structure factor $\langle \mathbf{S}_q \cdot \mathbf{S}_{-q} \rangle$. (b,e) Real-space spin-spin correlation function $\langle \mathbf{S}_{L/2} \cdot \mathbf{S}_\ell \rangle$, (c,f) nearest-neighbour chirality correlation function $\langle \kappa_{L/2}^1 \cdot \kappa_\ell^1 \rangle$. Top row [panels (a-c)] represents results for filling $n_K = 1/2$, while bottom row [panels (d-f)] depicts results for $n_K = 1/3$. All results calculated using DMRG and the generalized Kondo-Heisenberg model on $L = 48$ sites.

els is $n_K = 3 - n_H$.

In this work, we primarily reach our conclusions on the base of the gKH Hamiltonian, Eq. (2). However, in the Supplementary Notes we reproduce the main findings with the full two-orbital Hubbard model Eq. (1). All Hamiltonians are diagonalized via the density matrix renormalization group (DMRG) method where the dynamical correlation functions are obtained using the dynamical-DMRG technique (see Methods section). Open boundary conditions are assumed.

Let us briefly describe the several magnetic phases of the OSMP, where the novel phase reported here is in yellow in Fig. 1(c). For details we refer the interested reader to Ref. 24. In Fig. 1(c) we present a sketch of the interaction-filling (U - $n_{H/K}$) phase diagram: (i) At $U < W$ the ground state is a paramagnetic metal. (ii) For $U \gtrsim W$ the system enters OSMP with coexisting metallic and Mott-insulating bands. (iii) For sufficiently large values of interaction $U \gg W$ the system is in a FM state for all fillings. (iv) When $U \sim \mathcal{O}(W)$, namely when all energy scales compete, the system is primarily in the so-called block-magnetic state. Depending on the filling of the itinerant band, the spins form various sizes of AFM-coupled FM spin islands. An important special case, found experimentally in BaFe_2Se_3 , is the $n_K = 1/2$ ($n_H = 3/2 + 1$) filling where spins form a $\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\downarrow\downarrow$ -pattern along the legs, the $\pi/2$ -block magnetic state.

In between the block and FM phases, we discovered a novel region where the spin static structure factor $S(q) = \langle \mathbf{S}_q \cdot \mathbf{S}_{-q} \rangle$ [$\mathbf{S}_q = (1/\sqrt{L}) \sum_\ell \exp(-iq\ell) \mathbf{S}_\ell$ with $\mathbf{S}_\ell = \sum_\gamma \mathbf{S}_{\gamma,\ell}$] have its maximum q_{\max} at incommensu-

rate values of the wavevector q . In Figs. 2(a) and (d) we explicitly show the interaction U dependence of $S(q)$ in this region (see also Ref. 24) for $n_K = 1/2$ and $n_K = 1/3$, respectively. It is evident from the presented results that the maximum of the spin structure factor continuously changes with interaction U interpolating between block-magnetism at $U \sim W$ and ferromagnetic (FM) state at $U \gg W$ (e.g., at $n_K = 0.5$, from $q_{\max} = \pi/2$ to $q_{\max} = 0$). The real-space spin-spin correlation functions $\langle \mathbf{S}_{L/2} \cdot \mathbf{S}_\ell \rangle$ [see Fig. 2(b,e)] reveal an oscillatory structure throughout the chain, with period $\theta = q_{\max}$, decaying in amplitude at large spatial separations within the incommensurate region in the phase diagram. This behaviour may indicate spin-density wave or spiral order of the spin correlations. The competing interactions present in OSMP systems, and the location of this phase sandwiched between block and FM states, can potentially lead to a novel type of spiral state, as shown below. It is worth noting that θ represents an *average pitch* angle within the system. E.g., for the case of perfect $\pi/2$ -block $\uparrow\uparrow\downarrow\downarrow\uparrow\uparrow\downarrow\downarrow$ at $U \simeq W$ one gets π -angel (0-angel) between (within) the blocks.

Block spirals. To better investigate the magnetic structure of the novel spiral within the OSMP we will focus on the *chirality* correlation function $\langle \kappa_\ell^d \cdot \kappa_m^d \rangle$ [2, 3, 37, 38], where

$$\kappa_\ell^d = \mathbf{S}_\ell \times \mathbf{S}_{\ell+d}, \quad (3)$$

represents the vector product of two spin operators separated by a distance d , and consequently the angle be-




Figure 3. Chiral correlation function $\langle \kappa_\ell^d \cdot \kappa_m^d \rangle$. (a) Dependence of chirality $\kappa_\ell^d = \mathbf{S}_\ell \times \mathbf{S}_{\ell+d}$ on distance d between spins. Results calculated using the generalized Kondo-Heisenberg, $L = 48$, $n_K = 1/2$, and $U/W = 2.0$. Panel (b): same as (a) but for $n_K = 1/3$ and $U/W = 1.2$. In both panels: arrows represent schematics of the block order for given filling.

tween them. We stress that in the following we consider the total spin at each site $\mathbf{S}_\ell = \sum_\gamma \mathbf{S}_{\gamma,\ell}$. In the generic case of AFM- or FM-order, and also in the OSMF block-phase, the chirality correlation function vanishes, $\langle \kappa_\ell^d \cdot \kappa_m^d \rangle = 0$, since the spins are collinear. On the other hand, consecutive $\langle \kappa_\ell^d \cdot \kappa_m^d \rangle \neq 0$ indicate a nontrivial spiral order.

In Fig. 2(c,f) we present the interaction U dependence of the nearest-neighbour, $d = 1$, chirality correlation function $\langle \kappa_\ell^1 \cdot \kappa_m^1 \rangle$ of the gKH model. It is evident from the presented results that between the block and FM phases the chirality acquires finite values. In addition, the spatial structure of $\langle \kappa_\ell^1 \cdot \kappa_m^1 \rangle$ displays a clear zig-zag-like pattern. To better investigate the spiral internal structure, in Fig. 3 we present the dependence of the chirality correlation on the distance d between spins. In Supplementary Note 1 we provide the full interaction U dependence of the next-nearest-neighbour, $d = 2$, chirality correlation function. Here, as illustration we will focus on the representative cases of $U/W = 2.0$ for $n_K = 1/2$ and $U/W = 1.2$ for $n_K = 1/3$.

As already mentioned, the nearest-neighbour ($d = 1$) chiral correlation for both considered fillings contains additional patterns modulating the usual decay. Specifically, for $n_K = 1/2$ ($n_K = 1/3$) the correlation function “oscillates” every two (three) sites. Interestingly, these patterns change their nature when the next-nearest neighbour ($d = 2$) chirality is considered: (i) the values




Figure 4. Phase diagram varying the interaction U/W . Presented are: (i) maximum of static spin structure factor q_{\max} (squares), (ii) nearest- ($d = 1$) and next-nearest ($d = 2$) neighbour chirality $\tilde{\kappa}_{L/3}^d$ (open and filled circles, respectively), and (iii) dimer correlation $D_{\pi/2}$ (diamonds). Results calculated using the generalized Kondo-Heisenberg model, $L = 48$ sites, and $n_K = 1/2$.

of these chiralities increase $\langle \kappa_\ell^2 \cdot \kappa_m^2 \rangle > \langle \kappa_\ell^1 \cdot \kappa_m^1 \rangle$, and (ii) for the case of $n_K = 1/2$ the κ -correlation is now a smooth function of distance, while $n_K = 1/3$ still exhibits some three-site oscillations. Investigating the next-next-nearest neighbour case, $d = 3$, gives additional information. While for the $n_K = 1/2$ filling the $d = 3$ correlations are smaller than $d = 1$ and $d = 2$, for $n_K = 1/3$ they are larger and (as for $d = 2$ at $n_K = 1/2$) they are now a smooth function of distance. This seemingly erratic behaviour of $\langle \kappa_\ell^d \cdot \kappa_m^d \rangle$ correlations varying d cannot be simply explained by a mere uniform change of the pitch angle θ . The latter changes only with the interaction $\theta = \theta(U)$. On the other hand, as evident from Figs. 2(c,f), the internal structure of $\langle \kappa_\ell^1 \cdot \kappa_m^1 \rangle$ depends only on the electronic filling n_K (see also Supplementary Note 1 for $d = 2$ results).

To better explain this behaviour let us focus on the dimer correlation defined [39] as

$$D_{\pi/2} = \frac{2}{L} \sum_{\ell=L/4}^{3L/4} (-1)^{\ell-1} \langle \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+1} \rangle. \quad (4)$$

The above operator compares the number of FM and AFM bonds in the bulk system (Supplementary Note 2 contains the full real-space dependence of $\langle \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+1} \rangle$). For true FM or AFM ordered states, $|\uparrow\uparrow\uparrow\uparrow \dots\rangle$ or $|\uparrow\downarrow\uparrow\downarrow \dots\rangle$ respectively, each nearest-neighbour bond has the same sign: positive for FM $\uparrow\uparrow$ and negative for AFM $\uparrow\downarrow$. Consequently, $D_{\pi/2} = 0$. On the other hand, in the $\pi/2$ -block state, $|\uparrow\uparrow\downarrow\downarrow \dots\rangle$, the FM and AFM bonds alter in staggered fashion rendering $D_{\pi/2} \neq 0$. In Fig. 4 we present the interaction U dependence of $D_{\pi/2}$ for $n_K = 1/2$. Furthermore, in the same figure we present also the wavevector where the static structure factor is maximized q_{\max} , and the value of the $\tilde{\kappa}_{L/3}^d = \langle \kappa_\ell^d \cdot \kappa_m^d \rangle_{|\ell-m|=L/3}$ correla-

tor for $d = 1$ and $d = 2$. Starting in the paramagnet at small U both $D_{\pi/2}$ and $\tilde{\kappa}_{L/3}^d$ vanish, with $q_{\max} = \pi$ just depicting the usual short-range staggered correlations of weak- U physics. In the opposite limit of strong interaction, $U \gg W$, $D_{\pi/2} = \tilde{\kappa}_{L/3}^d = 0$ as well, consequence of a “simple” FM-state with $q_{\max} = 0$. In the most interesting case of competing interaction $U \sim W$, the dimer correlation $D_{\pi/2}$ acquires a finite value maximized at $U \simeq W$. The latter reflects a *perfect* $\pi/2$ -block magnetic state. Interestingly, one can observe a continuous transition of $D_{\pi/2}$ between the block- and FM-phases in the region where a finite chirality $\tilde{\kappa}_{L/3}^d \neq 0$ was found and where q_{\max} takes incommensurate values.

On the basis of the above results, a coherent picture emerges explaining the nature of the magnetic state between the block and FM limits. Consider first filling $n_K = 1/2$. At $U \simeq W$ the ground state is a block-magnetic phase, where two-site FM islands (blocks) are AFM coupled. Increasing the interaction U , the spins start to rotate w.r.t. each other, inducing finite $\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_m^d \rangle$ correlations. Remarkably, during the rotation the overall *FM islands*-nature of the state is preserved, yielding a finite $D_{\pi/2} \neq 0$ all the way to the FM-state at $U \gg W$. Such an unexpected scenario is also encoded in the inequalities $\tilde{\kappa}_{L/3}^2 > \tilde{\kappa}_{L/3}^1$ and $\tilde{\kappa}_{L/3}^2 > \tilde{\kappa}_{L/3}^3$ observed in Fig. 3(a). This is qualitatively different from a standard spiral state where the spin rotates from site to site [top sketch in Fig. 1(a)]. In our case, instead, the spiral is made of individual blocks, and it is the entire block that rotates from block-to-block [middle sketch in Fig. 1(a)].

Such a novel block-spiral state can also be observed at filling $n_K = 1/3$. Here, $\pi/3$ -blocks of three sites $\uparrow\uparrow\downarrow\downarrow$ develop a finite dimer correlation of the form $D_{\pi/3} \propto \sum_\ell f(\ell) \langle \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+1} \rangle$, where $f(\ell)$ accounts for the specific form of the bond sign pattern, i.e. $\{1, 1, -1, 1, 1, -1, \dots\}$ (see Supplementary Notes 2). A finite $D_{\pi/3}$ together with $\tilde{\kappa}_{L/3}^3 > \tilde{\kappa}_{L/3}^2 > \tilde{\kappa}_{L/3}^1$ is compatible with a spiral state of rotating three-site blocks [see bottom sketch in Fig. 1(a)].

Finally, let us comment on the finite-size dependence of our findings. Analysis of system sizes up to $L = 96$ sites (see Supplementary Note 3) indicates that the discussed block-spiral states display short-ranged order but with a robust correlation length of $\xi \sim 15$ sites, where $\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_{\ell+x}^d \rangle \propto \exp(-x/\xi)$. However, it was argued [2, 40] for the case of the FM long-range Heisenberg model that realistic small SU(2)-breaking anisotropies, often present in real materials, can induce true (quasi-)long-range order in a spiral state. Also, such an anisotropy will “choose” the plane of rotation of the spiral, i.e. in-plane or out-of-plane with regards to the chain direction. As we will argue in the next section, (frustrated) long-range Heisenberg Hamiltonians can (at least qualitatively) capture the main physics of the block-spiral unveiled here.

We emphasize that the same conclusions are reached in the multi-orbital Hubbard model, although because

the effort is much more computationally demanding it was limited to special cases. In Supplementary Note 4 we present results for $\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_m^d \rangle$ obtained with the full two-orbital Hubbard model (1) and also the incommensurability of $S(q)$ for a three-orbital Hubbard model.

Discussion

A distinctive feature of the OSMP is the coexistence of localized electrons (spins in an insulating band) and itinerant electrons (a metallic band). As shown below, the new block-spiral state discussed here is a consequence of the nontrivial interplay between localization and itinerancy.

In the block-magnetic phase at $U \simeq W$ it was previously shown [36] – for the three-orbital Hubbard model – that the density of states (DOS) at the Fermi level ϵ_F is reduced, indicating a pseudogap-like behaviour. Our calculations of the single-particle spectral function $A(q, \omega)$ and $\text{DOS}(\omega)$ (see Methods section) using the gKH model are presented in Fig. 5(a) and confirm this picture. They also show the strength of our effective Hamiltonian. We want to reiterate here that although the system is overall metallic in nature, the band structure is vastly different from the simple cosine-like result of $U \rightarrow 0$. Distinctive features in $A(q, \omega)$ at the Fermi vector k_F , and a large renormalization of the overall band structure at higher energies, indicate a complex interplay between various degrees of freedom and energy scales.

Upon increasing the interaction U and entering the block-spiral region, $A(q, \omega)$ changes drastically. In Fig. 5(b) we show representative results for a $\theta/\pi \simeq 0.3$ block-spiral state at $U/W = 2$ and $n_K = 1/2$. Two conclusions are directly evident from the presented results: (1) the pseudogap at ϵ_F is closed, but some additional gaps at higher energies opened. (2) $A(q, \omega)$ in the vicinity of the Fermi level, $\omega \sim \epsilon_F$, develops two bands, intersecting at the $q = 0$ and $q = \pi$ points, with maximum at $q \simeq \theta/2$. The bands represent two *quasiparticles*: “left”- and “right”-movers reflecting the two possible rotations of the spirals. It is obvious from the above results that the quasiparticles break parity symmetry, i.e., going from $k \rightarrow -k$ momentum changes the quasiparticle character, as expected for a spiral state. Somewhat surprisingly, $A(k, \omega)$ does not show any gap as would typically be associated with the finite dimerization D_α that we observe. However, it should be noted that for quantum localized $S = 1/2$ spins the quarter-filling ($n_K = 0.5$) implies a filling of $2/5$ of the lower Kondo band (due to energy difference between local Kondo singlets and triplets). The dimerization gap expected at $\pi/2$ would thus open away from the Fermi level and would consequently not confer substantial energy gain to the electrons. Similarly, no dimerization gap is found in the $n_K \simeq 0.33$ case, i.e., the $\pi/3$ -block spiral (not presented). We thus conclude that quantum fluctuations of the localized and itinerant




Figure 5. Spectral function $A(q, \omega)$. (a) Block-phase ($n_K = 1/2, U/W = 1.0$) and (b) block-spiral-phase ($n_K = 1/2, U/W = 2.0$) as calculated for the generalized Kondo-Heisenberg model using $L = 48$ sites. (c,d) The same results obtained with the two-orbital Hubbard model ($L = 48$). (c) Block-phase ($n_K = 2.50, U/W = 1.0$) and (d) block-spiral-phase ($n_K = 2.50, U/W = 2.0$). Horizontal (vertical) lines depict the Fermi level ϵ_F (Fermi wavevector k_F), while (i)PES stands for (inverse-) photoemission spectroscopy. Right panels: density of states $\text{DOS}(\omega) = \sum_q A(q, \omega)$.

spins are here strong enough to suppress the dimerization gap. The above conclusions can be reached from results obtained with the full two-orbital Hubbard model [see Figs. 5(c,d)]. Note that due to the particle-hole symmetry of the gKH model ($n_K \equiv n_H - 1 \equiv 3 - n_H$, where n_K and n_H stand for occupation in the gKH and two-orbital Hubbard models, respectively) the presented results for $n_K = 3 - n_H$ are “mirror” results of the two-orbital Hubbard model ($\omega \rightarrow -\omega$ and $k \rightarrow \pi - k$).

Since our system is in an overall metallic state, one could naively believe that the spin exchange J_r carried by the mobile electrons via Ruderman-Kittel-Kasuya-Yosida (RKKY) interaction could provide a suitable picture. As discussed below, this is not the case. It was previously shown [34] that the frustrated FM-AFM J_1 - J_2 Heisenberg model (with $|J_1| \sim J_2$) qualitatively captures the physics of the block-magnetic state. Here, in Fig. 6(a) we show that the spin structure factor $S(q)$ of the block-spiral state can be accurately described by a frustrated long-range Heisenberg Hamiltonian. Although this *phenomenological*

model is not derived here from the basic Hamiltonians, it accurately reproduces the interaction U dependence throughout the incommensurate region [e.g. compare Fig. 2(a) and Fig. 6(b)]. The intuitive understanding of the origin of the effective spin model is as follows: at $U \sim W$ the system is in a block-magnetic state with quasi-long-range (QLR) spin correlation of π/N_b nature, where N_b is the size of the block. This nontrivial effect of the interaction U , a metallic state coexisting with a (quasi)-ordered magnetic state, qualitatively modifies the nature of the RKKY interaction. We found excellent agreement between the gKH and Heisenberg models with $J_1 = -1, J_2 = 1/4, J_3 = 1/10$ (see also inset of Fig. 6 for the J_2 - J_3 dependence of q_{\max}). The latter yields $\alpha \simeq 2$ in $|J_r| \propto 1/r^\alpha$. As a consequence, the block-spiral magnetism is described by the class of Haldane-Shastry models [41, 42]. In fact, we speculate that due to the oscillating character of the electron mediated exchange couplings, the spin system which encompasses all phe-




Figure 6. Effective Heisenberg model. (a) Comparison of $S(q)$ between the generalized Kondo-Heisenberg (gKH) model in the spiral state ($L = 48, n_K = 1/2, U/W = 2.0$) and the long-range Heisenberg (lrH) model ($L = 48, J_2/|J_1| = 0.25, J_3/|J_1| = 0.1$). Note results are normalized by the magnetic moment $S^2 = \mathbf{S}(\mathbf{S} + 1)$, i.e., $S^2 = 1.375$ and $S^2 = 3/4$ for gKH and lrH, respectively. (b) Static spin structure factor $S(q)$ and (c) chirality correlation function $\langle \kappa_\ell^z \cdot \kappa_m^z \rangle$ of the J_1 - J_2 - J_3 Heisenberg model (with ferromagnetic $J_1 = -1$), calculated for $L = 120$ sites, $J_3/|J_1| = 0.1$, and various $J_2/|J_1| = 0.00, 0.05, \dots, 1.00$ (top to bottom). Inset of (a): position of the maximum of the static structure factor q_{\max} of the lrH model as function of $J_2/|J_1|$, for $J_3/|J_1| = 0.00, 0.05, 0.10$, and 0.15.

nomena is given by

$$H = \sum_{\ell, r} J_r \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+r},$$

$$\text{with } J_1 < 0, \quad J_2 > 0, \quad J_{r>2} = |J_1| \frac{(-1)^{r-1}}{r^\alpha}. \quad (5)$$

It was shown [43–45] that the above model have (for the zero magnetization sector, $S_{\text{tot}}^z = 0$) QLR ground

state with $\pi/2$ -correlations and also can support spiral states. Such frustrated long-range Hamiltonians are not suitable for direct DMRG calculations due to the area law of entanglement. Thus, alternatively in Supplementary Note 5 we show small system size Lanczos diagonalization results for Eq.(5). However, as presented in Fig. 6, the first three terms J_1 - J_2 - J_3 (which can still be computed accurately with DMRG) already give satisfactory results even for the chirality correlator [compare Fig. 2(c) and Fig. 6(c)].

Finally, it should be pointed out that the scenario (effective spin model) described above goes beyond the discussed $\pi/2$ -block case. Changes in the magnetization sector of the long-range FM Hamiltonian lead to modifications in the periodicity of the QLR order [46, 47]. As a consequence, there are only a few parameters in the effective spin model: the magnetization S_{tot}^z related to the filling in the original full multi-orbital Hamiltonian and $\{J_2, \alpha\} = f(U)$ which controls the long-range nature of the system.

Conclusion

In conclusion, we have identified a novel type of spiral spin order: block-spiral magnetism in which FM islands rigidly *rotate* with respect to each other. We wish to emphasize the crucial role of correlation in this phenomenon. Spiral states are usually a consequence of frustration (geometrical or induced by competing interactions) or by explicit symmetry breaking terms like DM couplings. On the contrary, in our system we have only nearest-neighbour interactions on a chain geometry, and the SU(2) symmetry is preserved. Instead, the novel block-spiral state reported here appears as an effect of competing interactions: the double-exchange like mechanism present in multi-orbital systems dominated by a robust Hund exchange, and the interaction U that governs superexchange tendencies. These nontrivial effects become apparent in the exotic effective (phenomenological) spin model: a long-range frustrated Heisenberg model. Within the latter the spin exchange decays slowly with distance, $\sim 1/r^2$, in contrast to the usual RKKY interaction which decays faster. Also, to our knowledge the multi-orbital system discussed in this work - as realized in iron-based compounds from the 123 family - could be one of the few, if not the only, realization of a Haldane-Shastry-like model with $\alpha = 2$.

Furthermore, due to properties unique to the OSMP, primarily the coexistence of metallic and insulating bands, the block-spiral state displays exotic phenomenon in the electronic degrees of freedom. For example, new quasiparticles appear due to the parity breaking of the spiral state (“left” and “right” movers). Similar physics can be found in systems with spin-orbit coupling [48, 49]. Here, again, this is an effect of competing energy scales. Another interesting possibility is the existence of mul-

tiferroic behaviour in our system. It is known [17–21] that in materials such as quasi-1D compounds LiCu_2O_2 , LiCuVO_2 or $\text{PbCuSO}_4(\text{OH})_2$, the spin spirals drive the system to ferroelectricity. Moreover, the phenomena described here, robust spiral magnetism without a charge gap, is at the heart of one of the proposed systems where topological Majorana phases can be induced [9–14].

Finally, let us comment on our results from the perspective of the real iron-based materials. As already mentioned, a nontrivial magnetic order, such as spirals, in the vicinity of high critical temperature superconductivity can lead to topological effects. This is the case of the 2D material $\text{FeTe}_{1-x}\text{Se}_x$ where zero-energy vortex bound states (Majorana fermions) have been reported [50–54]. Furthermore, similar to our findings, it was argued [55] that the frustrated magnetism of $\text{FeTe}_{1-x}\text{Se}_x$ can be captured by a long-range J_1 - J_2 - J_3 spin Hamiltonian. From this perspective, it seems appropriate to assume that the phenomena described in our work extends beyond 1D systems. Unfortunately, the lack of sufficiently reliable computational methods to treat 2D quantum models limits our understanding of multi-orbital effects in 2D. An intermediate promising route are the low-dimensional ladders from the family of 123 compounds where accurate DMRG calculations are possible. Early density functional theory and Hartree–Fock results suggest that the effects of correlations are important [35, 56] and that noncollinear magnetic order can develop in the ground-state [35]. The recent proposal of exploring ladder tellurides with a predicted higher value of U/W provides another avenue to consider [56]. As a consequence, we encourage crystal growers with expertise in iron-based materials to explore in detail the low-dimensional family of 123 compounds, including doped samples, because our results suggest that new and exotic physics may come to light.

Methods

DMRG method. The Hamiltonians discussed in this work were diagonalized using the density matrix renormalization group (DMRG) method [57–59] within the single-center site approach [60], where dynamical correlation functions are evaluated via the dynamical-DMRG [61–63] i.e. calculating spectral functions directly in frequency space with the correction-vector method [64] and Krylov decomposition [63]. The general premise of the method is to obtain a wave function that approximates the actual ground state. Within DMRG the latter is achieved by contracting a matrix-product state representing the ground state. The accuracy of the procedure can be quantified in terms of the discarded weight or truncation error which decreases with the size of the matrices or number of states kept (here we use up to 1200 states for all simulations).

Single-particle spectral function. The single particle

spectral function $A(q, \omega) = A^e(q, \omega) + A^h(q, \omega)$ is defined as

$$A^e(q, \omega) = -\frac{1}{\pi L} \sum_{\ell} e^{i\ell q} \text{Im} \langle \text{gs} | c_{\gamma, \ell}^{\dagger} \frac{1}{\omega^+ - H + \epsilon_0} c_{\gamma, L/2} | \text{gs} \rangle,$$

$$A^h(q, \omega) = -\frac{1}{\pi L} \sum_{\ell} e^{i\ell q} \text{Im} \langle \text{gs} | c_{\gamma, \ell} \frac{1}{\omega^+ + H + \epsilon_0} c_{\gamma, L/2}^{\dagger} | \text{gs} \rangle,$$

where $c_{\gamma, \ell} = c_{\gamma, \ell, \uparrow} + c_{\gamma, \ell, \downarrow}$, $\omega^+ = \omega + i\eta$, and $\eta = 2\Delta\omega$ with $\Delta\omega = 0.02$ as the frequency resolution. $A^e(q, \omega)$ [$A^h(q, \omega)$] represent retarded (electron photoemission) and advanced (hole inverse-photoemission) Green functions, respectively. The density of states can be calculated as $\text{DOS}(\omega) = \sum_q A(q, \omega)$ and similarly for the electron and hole parts.

Code availability. Computer codes used in this study are available at <https://github.com/dmrgPlusPlus/>.

Data Availability

The data that support the findings of this study are available from the corresponding author upon request.

-
- [1] Z. G. Soos, A. Parvej, and M. Kumar, *Numerical study of incommensurate and decoupled phases of spin-1/2 chains with isotropic exchange J_1 , J_2 between first and second neighbors*, *J. Phys.: Condens. Matter* **28** (2016) 175603.
 - [2] M. Sato, S. Furukawa, S. Onoda, and A. Furusaki, *Competing Phases in Spin-1/2 J_1 - J_2 Chain with Easy-Plane Anisotropy*, *Mod. Phys. Lett.* **25**, 901 (2011).
 - [3] I. P. McCulloch, R. Kube, M. Kurz, A. Kleine, U. Schollwöck, and A. K. Kolezhuk, *Vector chiral order in frustrated spin chains*, *Phys. Rev. B* **77**, 094404 (2008).
 - [4] M. J. P. Gingras and P. A. McClarty, *Quantum spin ice: a search for gapless quantum spin liquids in pyrochlore magnets*, *Rep. Prog. Phys.* **77**, 056501 (2014).
 - [5] N. Nagaosa and Y. Tokura, *Topological properties and dynamics of magnetic skyrmions*, *Nat. Nanotechnol.* **8**, 899 (2013).
 - [6] H. Tsunetsugu, Y. Hatsugai, K. Ueda, and M. Sigrist, *Spin-liquid ground state of the half-filled Kondo lattice in one dimension*, *Phys. Rev. B* **46**, 3175 (1992).
 - [7] L. Balents, *Spin liquids in frustrated magnets*, *Nature* **464**, 119 (2010).
 - [8] I. Kimchi, A. Nahum, and T. Senthil, *Valence Bonds in Random Quantum Magnets: Theory and Application to YbMgGaO_4* , *Phys. Rev. X* **8**, 031028 (2018).
 - [9] S. Nadj-Perge, I. K. Drozdov, J. Li, H. Chen, S. Jeon, J. Seo, A. H. MacDonald, B. A. Bernevig, Ali Yazdani, *Observation of Majorana fermions in ferromagnetic atomic chains on a superconductor*, *Science* **346**, 602 (2014).
 - [10] Y. Oreg, G. Refael, and F. von Oppen, *Helical Liquids and Majorana Bound States in Quantum Wires*, *Phys. Rev. Lett.* **105**, 177002 (2010).
 - [11] B. Braunecker and P. Simon, *Interplay between Classical Magnetic Moments and Superconductivity in Quantum*

- One-Dimensional Conductors: Toward a Self-Sustained Topological Majorana Phase*, *Phys. Rev. Lett.* **111**, 147202 (2013).
- [12] J. Klinovaja, P. Stano, A. Yazdani, and D. Loss, *Topological Superconductivity and Majorana Fermions in RKKY Systems*, *Phys. Rev. Lett.* **111**, 186805 (2013).
- [13] M. Schechter, M. S. Rudner, and K. Flensberg, *Spin-Lattice Order in One-Dimensional Conductors: Beyond the RKKY Effect*, *Phys. Rev. Lett.* **114**, 247205 (2015).
- [14] M. Steinbrecher, R. Rausch, K. T. That, J. Hermenau, A. A. Khajetoorians, M. Potthoff, R. Wiesendanger, and J. Wiebe, *Non-collinear spin states in bottom-up fabricated atomic chains*, *Nat. Commun.* **9**, 2853 (2018).
- [15] S. Dong, R. Yu, S. Yunoki, J.-M. Liu, and E. Dagotto, *Origin of multiferroic spiral spin order in the RMnO₃ perovskites*, *Phys. Rev. B* **78**, 155121 (2008).
- [16] I. A. Sergienko and E. Dagotto, *Role of the Dzyaloshinskii-Moriya interaction in multiferroic perovskites*, *Phys. Rev. B* **73**, 094434 (2006).
- [17] S. Seki, Y. Yamasaki, M. Soda, M. Matsuura, K. Hirota, and Y. Tokura, *Correlation between Spin Helicity and an Electric Polarization Vector in Quantum-Spin Chain Magnet LiCu₂O₂*, *Phys. Rev. Lett.* **100**, 127201 (2008).
- [18] S. Furukawa, M. Sato, and S. Onoda, *Chiral Order and Electromagnetic Dynamics in One-Dimensional Multiferroic Cuprates*, *Phys. Rev. Lett.* **105**, 257205 (2010).
- [19] Y. Tokura and S. Seki, *Multiferroics with Spiral Spin Orders*, *Adv. Mater.* **22**, 1554 (2010).
- [20] K. Yu. Povarov, Y. Feng, and A. Zheludev, *Multiferroic phases of the frustrated quantum spin-chain compound linarite*, *Phys. Rev. B* **94**, 214409 (2016).
- [21] A. Scaramucci, H. Shinaoka, M. V. Mostovoy, M. Müller, C. Mudry, M. Troyer, and N. A. Spaldin, *Multiferroic Magnetic Spirals Induced by Random Magnetic Exchanges*, *Phys. Rev. X* **8**, 011005 (2018).
- [22] J. M. Caron, J. R. Neilson, D. C. Miller, K. Arpino, A. Lobet, and T. M. McQueen, *Orbital-selective magnetism in the spin-ladder iron selenides Ba_{1-x}K_xFe₂Se₃*, *Phys. Rev. B* **85**, 180405(R) (2012).
- [23] L. de' Medici, S. R. Hassan, M. Capone, and X. Dai, *Orbital-Selective Mott Transition out of Band Degeneracy Lifting*, *Phys. Rev. Lett.* **102**, 126401 (2003).
- [24] J. Herbrych, J. Heverhagen, N. D. Patel, G. Alvarez, M. Daghofer, A. Moreo, E. Dagotto, *Novel Magnetic Block States in Low-Dimensional Iron-Based Superconductors*, *Phys. Rev. Lett.* **123**, 027203 (2019).
- [25] D. J. García, K. Hallberg, C. D. Batista, M. Avignon, and B. Alascio, *New Type of Charge and Magnetic Order in the Ferromagnetic Kondo Lattice*, *Phys. Rev. Lett.* **85**, 3720 (2001).
- [26] D. J. García, K. Hallberg, B. Alascio, and M. Avignon, *Spin Order in One-Dimensional Kondo and Hund Lattices*, *Phys. Rev. Lett.* **93**, 177204 (2004).
- [27] Q. Luo, K. Foyevtsova, G. D. Samolyuk, F. Reboredo, and E. Dagotto, *Magnetic states of the five-orbital Hubbard model for one-dimensional iron-based superconductors*, *Phys. Rev. B* **90**, 035128 (2014).
- [28] J. Rincón, A. Moreo, G. Alvarez, and E. Dagotto, *Exotic Magnetic Order in the Orbital-Selective Mott Regime of Multiorbital Systems*, *Phys. Rev. Lett.* **112**, 106405 (2014).
- [29] S. Wu, J. Yin, T. Smart, A. Acharya, C. L. Bull, N. Funnell, T. R. Forrest, G. Simutis, R. Khasanov, S. K. Lewin, M. Wang, B. A. Frandsen, R. Jeanloz, and R. J. Birgeneau, *Robust block magnetism in the spin ladder compound BaFe₂Se₃ under hydrostatic pressure*, arXiv:1909.06498 (2019).
- [30] M. Mourigal, S. Wu, M. B. Stone, J. R. Neilson, J. M. Caron, T. M. McQueen, and C. L. Broholm, *Block Magnetic Excitations in the Orbitaly Selective Mott Insulator BaFe₂Se₃*, *Phys. Rev. Lett.* **115**, 047401 (2015).
- [31] H. Takahashi, A. Sugimoto, Y. Nambu, T. Yamauchi, Y. Hirata, T. Kawakami, M. Avdeev, K. Matsubayashi, F. Du, C. Kawashima, H. Soeda, S. Nakano, Y. Uwatoko, Y. Ueda, T. J. Sato, and K. Ohgushi, *Pressure-induced superconductivity in the iron-based ladder material BaFe₂S₃*, *Nat. Mat.* **14**, 1008 (2015).
- [32] M. Wang, M. Yi, S. Jin, H. Jiang, Y. Song, H. Luo, A. D. Christianson, C. de la Cruz, E. Bourret-Courchesne, D.-X. Yao, D. H. Lee, and R. J. Birgeneau, *Spin waves and magnetic exchange interactions in the spin-ladder compound RbFe₂Se₃*, *Phys. Rev. B* **94**, 041111(R) (2016).
- [33] M. Wang, S. J. Jin, M. Yi, Y. Song, H. C. Jiang, W. L. Zhang, H. L. Sun, H. Q. Luo, A. D. Christianson, E. Bourret-Courchesne, D. H. Lee, D.-X. Yao, and R. J. Birgeneau, *Strong ferromagnetic exchange interaction under ambient pressure in BaFe₂S₃*, *Phys. Rev. B* **95**, 060502(R) (2016).
- [34] J. Herbrych, N. Kaushal, A. Nocera, G. Alvarez, A. Moreo, and E. Dagotto, *Spin dynamics of the block orbital-selective Mott phase*, *Nat. Commun.* **9**, 3736 (2018).
- [35] Q. Luo, A. Nicholson, J. Rincón, S. Liang, J. Riera, G. Alvarez, L. Wang, W. Ku, G. D. Samolyuk, A. Moreo, and E. Dagotto, *Magnetic states of the two-leg-ladder alkali metal iron selenides AFe₂Se₃*, *Phys. Rev. B* **87**, 024404 (2013).
- [36] N. D. Patel, A. Nocera, G. Alvarez, A. Moreo, S. Johnston, and E. Dagotto, *Fingerprints of an orbital-selective Mott phase in the block magnetic state of BaFe₂Se₃ ladders*, *Commun. Phys.* **2**, 64 (2019).
- [37] T. Hikihara, M. Kaburagi, H. Kawamura, and T. Tonegawa, *Ground State Phase Diagram of Frustrated S = 1 XXZ chains: Chiral Ordered Phases*, *J. Phys. Soc. Jpn.* **69**, 259 (2000).
- [38] T. Hikihara, M. Kaburagi, H. Kawamura, *Ground-state phase diagrams of frustrated spin-S XXZ chains: Chiral ordered phases*, *Phys. Rev. B* **63**, 174430 (2001).
- [39] J. C. Xavier, R. G. Pereira, E. Miranda, and I. Affleck, *Dimerization Induced by the RKKY Interaction*, *Phys. Rev. Lett.* **90**, 247204 (2003).
- [40] M. F. Maghrebi, Z.-X. Gong, and A. V. Gorshkov, *Continuous Symmetry Breaking in 1D Long-Range Interacting Quantum Systems*, *Phys. Rev. Lett.* **119**, 023001 (2017).
- [41] B. S. Shastry, *Exact solution of an S = 1/2 Heisenberg antiferromagnetic chain with long-ranged interactions*, *Phys. Rev. Lett.* **60**, 639 (1988).
- [42] F. D. M. Haldane, *Exact Jastrow-Gutzwiller resonating-valence-bond ground state of the spin-1/2 antiferromagnetic Heisenberg chain with 1/r² exchange*, *Phys. Rev. Lett.* **60**, 635 (1988).
- [43] A. W. Sandvik, *Ground States of a Frustrated Quantum Spin Chain with Long-Range Interactions*, *Phys. Rev. Lett.* **104**, 137204 (2010).
- [44] N. Laflorencie, I. Affleck, and M. Berciu, *Critical phenomena and quantum phase transition in long range Heisenberg antiferromagnetic chains*, *J. Stat. Mech.*

- P12001 (2005).
- [45] Z.-H. Li and A.-M. Wang, *Matrix product state approach to a frustrated spin chain with long-range interactions*, *Phys. Rev. B* **91**, 235110 (2015).
 - [46] H. Ueda and S. Onoda, *Vector-spin-chirality order in a dimerized frustrated spin-1/2 chain*, *Phys. Rev. B* **89**, 024407 (2014).
 - [47] H. Onishi, *Magnetic Excitations of Spin Nematic State in Frustrated Ferromagnetic Chain*, *J. Phys. Soc. Jpn.* **84**, 083702 (2015).
 - [48] G. W. Winkler, M. Ganahl, D. Schuricht, H. G. Evertz and S. Andergassen, *Interaction effects in a microscopic quantum wire model with strong spinorbit interaction*, *New J. Phys.* **19**, 063009, (2017).
 - [49] X. Li, E. Zhao, and W. V. Liu, *Topological states in a ladder-like optical lattice containing ultracold atoms in higher orbital bands*, *Nat. Commun.* **4**, 1523 (2012).
 - [50] P. Zhang, K. Yaji, T. Hashimoto, Y. Ota, T. Kondo, K. Okazaki, Z. Wang, J. Wen, G. D. Gu, H. Ding, S. Shin, *Observation of topological superconductivity on the surface of an iron-based superconductor*, *Science* **360**, 182 (2018).
 - [51] D. Wang, L. Kong, P. Fan, H. Chen, S. Zhu, W. Liu, L. Cao, Y. Sun, S. Du, J. Schneeloch, R. Zhong, G. Gu, L. Fu, H. Ding, H.-J. Gao, *Evidence for Majorana bound states in an iron-based superconductor*, *Science* **362**, 333 (2018)
 - [52] P. Zhang *et al.*, *Multiple topological states in iron-based superconductors*, *Nat. Phys.* **15**, 41 (2018).
 - [53] T. Machida, Y. Sun, S. Pyon, S. Takeda, Y. Kohsaka, T. Hanaguri, T. Sasagawa, and T. Tamegai, *Zero-energy vortex bound state in the superconducting topological surface state of Fe(Se,Te)*, *Nat. Mat.* **18**, 811 (2019).
 - [54] R.-X. Zhang, W. S. Cole, and S. Das Sarma, *Helical Hinge Majorana Modes in Iron-Based Superconductors*, *Phys. Rev. Lett.* **122**, 187001 (2019).
 - [55] H. Ruiz, Y. Wang, B. Moritz, A. Baum, R. Hackl, and T. P. Devereaux *Frustrated magnetism from local moments in FeSe*, *Phys. Rev. B* **99**, 125130 (2019).
 - [56] Y. Zhang, L.-F. Lin, A. Moreo, S. Dong, and E. Dagotto, *Magnetic states of iron-based two-leg ladder tellurides*, [arXiv:1908.11765](https://arxiv.org/abs/1908.11765) (2019).
 - [57] S. R. White, *Density matrix formulation for quantum renormalization groups*, *Phys. Rev. Lett.* **69**, 2863 (1992).
 - [58] U. Schollwöck, *The density-matrix renormalization group*, *Rev. Mod. Phys.* **77**, 259 (2005).
 - [59] G. Alvarez, *The Density Matrix Renormalization Group for Strongly Correlated Electron Systems: A Generic Implementation*, *Comput. Phys. Commun.* **180**, 1572 (2009).
 - [60] S. R. White, *Density matrix renormalization group algorithms with a single center site*, *Phys. Rev. B* **72**, 180403 (2005).
 - [61] E. Jeckelmann, *Dynamical density-matrix renormalization-group method*, *Phys. Rev. B* **66**, 045114 (2002).
 - [62] H. Benthien and E. Jackelmann, *Spin and charge dynamics of the one-dimensional extended Hubbard model*, *Phys. Rev. B* **75**, 205128 (2007).
 - [63] A. Nocera and G. Alvarez, *Spectral functions with the density matrix renormalization group: Krylov-space approach for correction vectors*, *Phys. Rev. E* **94**, 053308 (2016).
 - [64] T. D. Kühner and S. R. White, *Dynamical correlation*

functions using the density matrix renormalization group, *Phys. Rev. B* **60**, 335 (1999).

Acknowledgments

We thank M. L. Baez, C. Batista, and M. Mierzejewski for fruitful discussions. J. Herbrych, A. Moreo, and E. Dagotto were supported by the US Department of Energy (DOE), Office of Science, Basic Energy Sciences (BES), Materials Sciences and Engineering Division. In addition, J. Herbrych acknowledges grant support by the Polish National Agency of Academic Exchange (NAWA) under contract PPN/PPO/2018/1/00035. The work of G. Alvarez was supported by the Scientific Discovery through Advanced Computing (SciDAC) program funded by the U.S. DOE, Office of Science, Advanced Scientific Computer Research and Basic Energy Sciences, Division of Materials Science and Engineering. The development of the DMRG++ code by G. Alvarez was conducted at the Center for Nanophase Materials Science, sponsored by the Scientific User Facilities Division, BES, DOE, under contract with UT-Battelle. J. Heverhagen and M. Daghofer were supported by the Deutsche Forschungsgemeinschaft, via the Emmy-Noether program (DA 1235/1-1) and FOR1807 (DA 1235/5-1) and by the state of Baden-Württemberg through bwHPC. Calculations have been carried out using resources provided by Wroclaw Centre for Networking and Supercomputing.

Author contributions

J. Herbrych and E. Dagotto planned the project. J. Herbrych and J. Heverhagen performed numerical calculations. G. Alvarez developed the DMRG++ computer program. A. Moreo and M. Daghofer developed the quasiparticle picture. J. Herbrych, G. Alvarez, A. Moreo, and E. Dagotto wrote the manuscript. All co-authors provided comments on the paper.

Additional information

Supplementary Information accompanies this paper.

SUPPLEMENTARY INFORMATION for:
Block-spiral magnetism: A New Type of Frustrated Order

by J. Herbrych, J. Heverhagen, G. Alvarez, M. Daghofer, A. Moreo, E. Dagotto

SUPPLEMENTARY NOTE 1
Chirality correlations

In Fig. S1 we present data that is complementary to that shown in Fig. 3 of the main text for the next-nearest neighbour, $d = 2$, chirality correlation function. These results are for the generalized Kondo-Heisenberg model and calculated for $n_K = 1/2$ [panel (a)] and $n_K = 1/3$ [panel (b)] (using $L = 48$ sites). As described in the main text, $\langle \kappa_\ell^2 \cdot \kappa_m^2 \rangle$ becomes a smooth function of distance for $n_K = 1/2$ for most considered values of the interaction U , particularly in the intermediate region where this correlation is robust, indicative of a spiral that preserves the building blocks of the two-down two-up state. On the other hand, the $n_K = 1/3$ case exhibits three-site oscillations in a “two-one” pattern even when this correlation is robust in value. This is compatible with the three-down three-up block state of this density.




Figure S1. Interaction U dependence of the next-nearest neighbour, $d = 2$, chirality correlation function $\langle \kappa_{L/2}^2 \cdot \kappa_\ell^2 \rangle$, using the generalized Kondo-Heisenberg model on $L = 48$ sites. Panel (a) are results for filling $n_K = 1/2$, while panel (b) depicts results for $n_K = 1/3$.

SUPPLEMENTARY NOTE 2
Dimer correlation

In Fig. S2 we present the bond order operator $\langle \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+1} \rangle$ throughout the chain investigated. As described in the main text, this local dimer correlatoin D_{order} allow us to visualize the number of AFM and FM bonds in the states investigated. It is evident from the results depicted in Figs. S2(c) and (d) that the block-magnetic order can be accurately identified by D_{order} .




Figure S2. Spatial dependence of the bond order operator $\langle \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+1} \rangle$ for various states: (a) FM state, (b) AFM state, (c) $\pi/2$ -block state, and (c) $\pi/3$ -block state. All results are calculated using the generalized Kondo-Heisenberg model on $L = 48$ sites.

SUPPLEMENTARY NOTE 3 Size dependence

In Figs. S3(a,b), we present the system-size dependence of the nearest- and next-nearest neighbour ($d = 1$ and $d = 2$, respectively) chirality correlation function $\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_m^d \rangle$ for systems up to $L = 96$ sites. Our results [see Fig. S3(c)] suggest that the $\boldsymbol{\kappa}$ -correlators decay exponentially with a robust correlation length $\xi \simeq 15$, where

$$\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_{\ell+x}^d \rangle \propto \exp(-x/\xi). \quad (\text{S1})$$




Figure S3. System-size dependence of the chirality correlation function. (a) Nearest- and (b) next-nearest neighbour chirality correlation $\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_m^d \rangle$ calculated for various system sizes $L = 24, \dots, 96$. We use the generalized Kondo-Heisenberg model for $n_K = 1/2$ and $U/W = 2.0$. (c) Log-plot illustrating the distance dependence of the chirality correlation $\langle \boldsymbol{\kappa}_\ell^d \cdot \boldsymbol{\kappa}_m^d \rangle$ function as calculated for $L = 96$. The solid line represents a fit to the function $f(x) \propto \exp(-x/\xi)$.

In Fig. S4 we present the system-size dependence of the static spin structure factor $S(q) = \langle \mathbf{S}_{-q} \cdot \mathbf{S}_q \rangle$. In agreement with results for the chirality correlation functions, we find indications of short-range order behaviour because the peak strength does not continue growing with increasing L , but instead converges to a fixed value. Note that spirals with longer wavelengths (as in the case presented in Fig. S4(a) for $n_K = 1/3$) have stronger finite-size effects: this is natural because here the elementary building blocks involve three spins, instead of only two at $n_K = 1/2$.




Figure S4. System-size dependence of the spin structure factor $S(q)$. Panel (a) presents data for $n_K = 1/3$ and $U/W = 1.2$ and (b) for $n_K = 1/2$ and $U/W = 2$. Shown are results for various system sizes $L = 24, \dots, 96$ using the generalized Kondo-Heisenberg model.

SUPPLEMENTARY NOTE 4 Multi-orbital results

In Fig. S5 we repeat the main findings of our work, namely the nearest- and next-nearest neighbour chirality correlation functions as shown for example in Fig. 3 of the main text, but now using the more complete, and more difficult, two-orbital Hubbard model,

$$\begin{aligned} H_H = & - \sum_{\gamma, \gamma', \ell, \sigma} t_{\gamma \gamma'} \left(c_{\gamma, \ell, \sigma}^\dagger c_{\gamma', \ell+1, \sigma} + \text{H.c.} \right) + \Delta \sum_{\ell} n_{1, \ell} \\ & + U \sum_{\gamma, \ell} n_{\gamma, \ell, \uparrow} n_{\gamma, \ell, \downarrow} + (U - 5J_H/2) \sum_{\ell} n_{0, \ell} n_{1, \ell} \\ & - 2J_H \sum_{\ell} \mathbf{S}_{0, \ell} \cdot \mathbf{S}_{1, \ell} + J_H \sum_{\ell} \left(P_{0, \ell}^\dagger P_{1, \ell} + \text{H.c.} \right) \quad (\text{S2}) \end{aligned}$$

In agreement with the results presented in Fig. 3 of the main text, the present results obtained with the multi-orbital system are fully consistent with the discussed block-spiral state displayed in the main text.

Let us now comment on other multi-band models. In particular, we will consider the following Hamiltonians closely related to the generalized Kondo-Heisenberg model:

- Kondo lattice (KL) — $U = K = 0$ in gKH,
- Kondo Heisenberg (KH) — $U = 0$ in gKH.

Furthermore, we will investigate the gKH model where only one system parameter (either the Hund exchange J_H or the interaction U) is changed. Finally, we will present results for the full three-orbital Hubbard model




Figure S5. Chiral correlation functions $\langle \kappa_\ell^d \cdot \kappa_m^d \rangle$ with $\kappa_\ell^d = \mathbf{S}_\ell \times \mathbf{S}_{\ell+d}$. (a) Dependence of these chiral correlation functions vs distance between spins. Results calculated for the two-orbital Hubbard model, $L = 48$, $n_H = 2.50$, and $U/W = 2.0$. Panel (b): same as (a) but for $n_H = 2.66$ and $U/W = 1.2$.

[28, 34, 36] which exhibits OSMP with one localized and two itinerant bands. In the latter, we use $t_{00} = t_{11} = -0.5$, $t_{22} = -0.15$, $t_{01} = t_{10} = 0$, $t_{02} = t_{20} = 0.1$ together with $\Delta_0 = -0.1$, $\Delta_1 = 0$, $\Delta_2 = 0.8$, all in units of eV. For all considered systems we fix the units to the kinetic energy bandwidth of gKH, i.e., $W = 2.1$ eV. Some additional comments are necessary: (i) it is important to note that our KL and KH models have a factor of 2 in front of the Hund exchange – remnant of the original derivation of gKH. (ii) For the KH model, we fixed the value of the spin exchange to $K = 4t_{11}^2/(2W) = 0.0214$. (iii) For the results of gKH with only one variable being changed at a time we fix the remaining parameters to the value corresponding to $U/W = 2$. (iv) For all Kondo-like models (KL, KH, gKH) we choose the $n_K = 1/2$ filling, namely $\pi/2$ -block magnetic states in block-phase. The latter is stabilized [28, 34] for $n_H = 4/3$ in the three-orbital Hubbard model which we use here.

A good indicator of the spiral state in our investigation is the incommensurability of the maximum q_{\max} of the static structure factor $S(q)$. In Fig. S6 we present the Hund exchange and interaction U dependence of q_{\max} for various models. For KL we reproduce results [26] where the spiral state is stabilized for $6 \lesssim J_H/t \lesssim 10$. Interestingly, introducing a finite exchange between localized spins $K \neq 0$ pushes the spirals to a larger range value of Hund exchange. Such a behaviour is an illustration of competing energy scales in our system: although the relatively small K naively appears irrelevant, i.e. $K/J_H \sim 0.01$, it has considerable effect on $S(q)$. The effect of finite interaction U/W also modifies the dependence of spirals on Hund exchange. In the latter the

relevant energy scale is reduced by a factor of ~ 4 .

Let us now comment on the effect of interaction U/W . As already shown in the main text, in the gKH model the block-spiral state can be found in the range $1.7 \lesssim U/W \lesssim 2.3$. The two-orbital Hubbard model yields the same results [24]. In the iron-based materials, for which our models are relevant, interaction U/W can be naively controlled by, e.g., pressure. Assuming that other system parameters are not affected by such an action, the spiral region can be enlarged. In Fig. S6(b) we show that changes only in the interaction U/W stabilizes block-spiral state in the range $1.2 \lesssim U/W \lesssim 2.5$. Finally, to test the robustness of our findings we investigated the three-orbital Hubbard model. As is evident from Fig. S6(b), between the block and FM phases there is a region of incommensurate value of $S(q)$ and finite dimerization $D_{\pi/2}$ (not presented). As a consequence, the block-spiral state is evidently robust against modifications of the system parameters and even the number or orbitals, and thus we argue that it is a robust property of the OSMP itself.




Figure S6. Dependence of the maximum q_{\max} of the static structure factor $S(q)$ on the value of (a) the Hund exchange J_H/W and (b) the interaction U/W . In (a) we present results for the Kondo lattice (KL, $U = K = 0$), Kondo-Heisenberg (KH, $U = 0$, $K = 0.0214$), and generalized Kondo-Heisenberg (gKH) models with parameters fixed (to the value given by $U/W = 2$) except the Hund exchange coupling. In (b) we present results for the full gKH, gKH with parameters fixed (to the value given by $U/W = 2$) except interaction, and three-orbital Hubbard model (see text for details). Colored areas represent regions where spiral states are found.

SUPPLEMENTARY NOTE 5

Frustrated long-range Heisenberg model

In the main text, we have shown that qualitatively the block-chirality can be described effectively by the J_1 - J_2 - J_3 Heisenberg model. It may be suspected that due to the oscillating nature of the mobile-electrons mediated spin-exchange, together with the long-range nature of spin correlations within OSMP, the generic effective spin-1/2 Hamiltonian that must be used to describe the discussed phenomena should be more extended and of the form

$$H = \sum_{\ell,r} J_r \mathbf{S}_\ell \cdot \mathbf{S}_{\ell+r}, \quad (\text{S3})$$

with $J_1 < 0$, $J_2 > 0$, and $J_{r>2} = |J_1| \frac{(-1)^{r-1}}{r^\alpha}$.

Such a frustrated long-range Hamiltonian is not suitable for DMRG-like approaches due to the well-known area law of entanglement which ‘‘controls’’ the accuracy of simulation. Thus, in order to gain understanding of the behaviour of the static structure factor $S(q)$ on such a model with extended interactions, we have used exact Lanczos diagonalization on a chain of $L = 20$ sites with OBC.

In Fig. S7 we present the dependence of the position of the peak q_{\max} on the J_2 -interaction and on α in the Eq. (S3) model. Panel (a) depicts results for overall magnetization $S_{\text{tot}}^z = 0$ while panel (b) for $S_{\text{tot}}^z = 1/4$. Depending on the magnetization, at large J_2 the system is dimerized with quasi-long range order (QLRO) of $q_{\max} = \pi/N_b$ nature [47], where N_b is the size of the block (e.g. $N_b = 2$ for $S_{\text{tot}}^z = 0$). On the other hand, in the $J_2 \rightarrow 0$ limit the usual FM-order is stabilized. In between those two phases, similarly as in the case of the gKH model, the maximum of the spin structure factor acquires incommensurate values. It is known [46] that within this region a spiral can develop. A generic phase diagram of such a scenario is presented in Fig. S7(c).

In summary, in this Note 5 we show that the successive FM-spiral-block transitions of the effective J_1 - J_2 - J_3 model described in the main text do not occur only in a narrow region of parameters. Here we have shown that a more generic model, with only two parameters $J_2/|J_1|$ and α , comfortably supports such a physics. For this reason, our conclusions do not require fine tuning of couplings but are broadly universal and should appear in many models, and hopefully also in associated real materials.




Figure S7. Position of the maximum in the spin structure factor, q_{\max} , for the frustrated long-range Heisenberg model described in this Note. In panel (a) we present results from magnetization $S_{\text{tot}}^z = 0$, while in (b) for $S_{\text{tot}}^z = 1/4$. Results were calculated using $L = 20$ sites and the Lanczos diagonalization. (c) Schematic phase diagram of the frustrated long-range Heisenberg model.